

General Information for 1999 corn rootworm insecticide test. Gerald Wilde, Kansas State University, Department of Entomology, Manhattan, KS.

Pest: Western Corn Rootworm, Diabrotica vergifera vergifera

Crop: Corn

Location: Manhattan, KS. Site 2

Planting Date: April 21

Herbicide: 4/21 Broadstrike 55 + Dual @ pt 2/acre + Atrex 90 @ 1 lb/acre.

5/13 - 2-4 - Do 2 pt/acre

Plot size: 1 row 30 ft. long, 4 replications, 30 inch row spacing

Experimental Design: Randomized Complete Block, 8 Treatments

Planting Information: Planted corn 1 2 inches in depth.

Field History: Corn 1997

Fertilization: 2/18-28-0-0 at 30 gals/acre

Application Information: Insecticides applied with v-belt seeder in T Band.

Evaluation: Corn rootworm larvae damage rating on 4 plants per plot on July 6, 1999.

Scale 1-6: 1=no damage; 6=severe damage.

Phytotoxicity: none noted

Rainfall and Temperature: enclosed

Soil: pH 5.2, OM 3.9%, sand 18%, silt 56%, clay 26%, CEC 27.1

Evaluation: Good rootworm pressure was present as noted by root ratings in the

untreated plots. Some treatments were more effective than others.

1999 corn rootworm insecticide test.

Gerald Wilde, Kansas State University, Department of Entomology, Manhattan, KS.

TRT	FORM	RATE ¹	PLACE ²	TIME ³	ROOT
1. Regent	4 SC	$\overline{0.24 \text{ fl.}}$ oz	ĪF	P	1.38 a
2. Aztec	2.1 G	6.7 oz	TB	P	1.56 abc
3. Aztec	2.1 G	6.7 oz	IF	P	1.44 ab
4. Aztec	2.1 B	6.7 oz	TB	P	1.56 abc
5. Aztec	2.1 B	6.7 oz	IF	P	1.81 abcd
6. Counter	20 CR	6 oz	TB	P	2.81 ghij
7. Capture	1.1 G	5.3 oz	TB	P	3.31 jkl
8. Capture	1.1 G	6.4 oz	TB	P	3.00 hijk
9. Capture	1.1 G	8 oz	TB	P	3.13 ijk
10. Capture	2 E	0.05 a.i.	TB	P	2.50 defghij
11. Capture	2 E	0.06 a.i.	TB	P	2.25 bcdefgh
12. Capture	2 E	0.075 a.i.	TB	P	2.31 cdefghi
13. Force	3 G	4 oz	IF	P	2.81 ghij
16. Lorsban	75 WG	1.3 a.i.	TB	P	1.88 abcef
17. Lorsban	75 WG	1.3 a.i	IF	P	1.69 abcd
18. Lorsban	15 G	8 oz	TB	P	2.69 efghij
19. Lorsban	15 G	8 oz	IF	P	3.13 ijk
20. Lorsban	4 E	1.3 a.i.	TB	P	2.44 defghi
21. Thimet	20 G	6 oz	TB	P	2.75 fghij
22. Fortress	5 G	3 oz	TB	P	2.13 abcdefg
23. Force	3 G	4 oz	TB	P	1.38 a
24. Force	WF 2.5	0.24 fl. oz	В	P	2.00 abcdefg
25. Force	WF 2.5	0.36 fl. oz	В	P	2.69 efghij
26. Force	WF 2.5	0.48 fl. oz	В	P	1.38 a
27. Force	WF 2.5	0.24 fl. oz	IF	P	1.94 abcdef
28. Force	WF 2.5	0.36 fl. oz	IF	P	2.13 abcdefg
29. Force	WF 2.5	0.48 fl. oz	IF	P	1.69 abcd
30. Lorsban	4 E	1.3 a.i.	IF	P	2.44 defghi
31. Furadan	4 E	1.0	В	P	4.00 lmn
32. Furadan	4 E	1.0	IF	P	3.75 klm
33. Untreated					4.82 n
34. Furadan	4 F	1.0	BR	C	2.13 abcdefg
36. Untreated					4.25 mn

¹ Oz. per 1000 row ft or lb. a.i./acre

² IF - infurrow, TB-T-Band, BR-Broadcast ³ P = Planting, C = Cultivation

Reference to specific products is provided solely for informational purposes. Experiments with pesticides on non-labeled crops or pests is part of the insecticide registration process, it does not imply endorsement or recommendation of non-labeled uses of pesticides by Kansas State University. All pesticide use must be consistent with current labels.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service It is the policy of Kansas State University Agricultural Experiment Station and Cooperative Extension Service that all persons shall have equal opportunity and access to its educational programs, services, activities, and materials without regard to race, color, religion, national origin, sex, age or disability. Kansas State University is an equal opportunity organization. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, Marc A. Johnson, Director.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service