



Sugarcane rootstock weevil control on sorghum 2004*

Abilene, Kansas

Investigator: Dr. Gerald Wilde

Crop 1: SORBI SORGHUM BICOLOR
 Variety: DK54
 Planting Date: 25/May/2004
 Planting Method: John Deere Planter
 Rate: 40000 P/A
 Depth: 2 IN

Application Date: 25/Jun/2004
 Time of Day: 10:00am
 Application Method: hand spray
 Application Timing: POSPOS
 Application Placement: INWHOR
 Air Temp., Unit: 80 F
 Crop 1 Code, Stage: SORBI whorl
 Stage Scale:
 Height, Unit: 1 FT

Rating Date 27/Jun/2004

No.	Treatment/ Product Name	Weevils per Plant
1	Lorsban @ 1.0	0.00 b
2	Warrior @ 0.03	0.25 b
3	Mustang @ 0.02	0.25 b
4	Baythroid @ 0.037	0.00 b
5	Furadan @ 0.5	0.00 b
6	Asana @ 0.03	0.50 b
7	Untreated	4.25 a

Means followed by same letter do not significantly differ ($P > .05$; PROC GLM; Mean comparison by LSD [SAS institute 2003])

*Revised Jan. 9, 2005 pes

Reference to specific products is provided solely for informational purposes. Experiments with pesticides on non-labeled crops or pests is part of the insecticide registration process, it does not imply endorsement or recommendation of non-labeled uses of pesticides by Kansas State University. All pesticide use must be consistent with current labels.

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, Fred Cholick, Director.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service