



Stem Borer Seed Treatment Beloit, Kansas 2003

Gerald Wilde, Kansas State University, Department of Entomology, Manhattan, KS

Pest: Soybean stem borer

Crop: Sunflower

Rating Data Type			
Trt No.	Treatment Name	Rate Unit	Percent Lodging
1	Untreated		22.75 a
2	Apron + Cruiser		30.25 a
3	Apron + Cruiser		23.25 a
4	Apron + Cruiser		31.25 a
LSD (P=.05)			8.255
Standard Deviation			5.161
CV			19.2
Grand Mean			26.88
Bartlett's X2			0.869
P(Bartlett's X2)			0.833
Replicate F			9.701
Replicate Prob(F)			0.0035
Treatment F			3.038
Treatment Prob(F)			0.0855

Means followed by same letter do not significantly differ (P=.05, Student-Newman-Keuls)

Mean comparisons performed only when AOV Treatment P(F) is significant at mean comparison OSL.

Reference to specific products is provided solely for informational purposes. Experiments with pesticides on non-labeled crops or pests is part of the insecticide registration process, it does not imply endorsement or recommendation of non-labeled uses of pesticides by Kansas State University. All pesticide use must be consistent with current labels.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, Fred A. Cholick, Director.