



White Grub Control – Corn (2005) – Field Test

Planting date: 04/22/05

Evaluation date: 05/16/05

Gerald E. Wilde - Kansas

Trt. No.	Treatment/ Product Name	plants/hill (Mean ± SE)
1	Untreated	0.40 ± 0.40d
2	Cruiser 5 FS @ 0.125 mg/seed	2.00 ± 0.00abc
3	A14115 @ 0.138 mg/seed + Cruiser 5 FS @ 1.125 mg/seed	1.20 ± 0.49cd
4	Cruiser 5 FS @ 0.25 mg/seed	2.00 ± 0.00abc
5	Cruiser 5 FS @ 1.125 mg/seed	2.00 ± 0.00abc
6	STP15199 @ 5 mg/seed + STP15201 @ 0.25 mg/seed	2.40 ± 0.24a
7	STP15142 2.0 gm/100 Kg seed + STP15281 @ 58.5 gm/100 Kg seed	2.20 ± 0.37ab
8	Poncho 1250 ST @ 1.25 mg/kernal	2.00 ± 0.32abc
9	Poncho 1250 ST @ 0.25 mg/kernal	1.40 ± 0.40bc

Means within a column followed by the same letter are not significantly different ($P > 0.05$; PROC GLM; Mean comparison by LSD [SAS Institute 2003]).

Reference to specific products is provided solely for informational purposes. Experiments with pesticides on non-labeled crops or pests is part of the insecticide registration process, it does not imply endorsement or recommendation of non-labeled uses of pesticides by Kansas State University. All pesticide use must be consistent with current labels.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, Fred A. Cholick, Director.