



Corn Rootworm Control in Field Corn with planting time treatments. Soil
Insecticide Test # 3, 2005

Gerald Wilde, Department of Entomology, Kansas State University

Pest:	Western Corn Rootworm, <i>Diabrotica virgifera virgifera</i>
Crop:	Corn, <i>Zea mays</i> L. 26 treatments
Location:	Manhattan, Kansas
Planting Date:	April 29, 2005
Plot Size:	1 rows, 25 ft
Experimental Design:	Randomized Complete Block; 4 Replications
Planting Information:	Granules applied with v-belt seeder. Corn planted 1-2 inch depth. Soil in good moist condition at planting, 30 inch rows, disc 2 inch just before planting. Infurrow treatments applied with seed..
Field History:	Corn 2004
Phytotoxicity:	none noted
Evaluation:	Evaluated on July 01, 2005; Evaluated 4 plants/replication, total 16 plants/treatment; Damage rating using 0-3 scale where 0 = no damage and 3 = 3 nodes destroyed

Soil Insecticide – Test 3 (North Farm, Manhattan)
 Corn Rootworm – Corn (2005)
 Planting date: April 29, 2005
 Evaluation date: July 01, 2005

Gerald Wilde, Department of Entomology, Kansas State University

Trt. No.	Treatment/ Product Name	Application	Root Rating (Mean ± SE)
1	Untreated Check	----	0.82 ± 0.19ab
2	Force 3G @ 4 OzPr/1000 row ft.	Band	0.16 ± 0.06def
3	Force 3G @ 4 OzPr/1000 row ft.	IF	0.07 ± 0.03f
4	A14776 @ 0.09 OzA/1000 row ft.	Band	0.39 ± 0.14cd
5	A14776 @ 0.12 OzA/1000 row ft.	Band	0.37 ± 0.14cde
6	A14776 @ 0.15 OzA/1000 row ft.	Band	0.11 ± 0.04ef
7	A14776 @ 0.09 OzA/1000 row ft.	IF	0.58 ± 0.24bc
8	A14776 @ 0.12 OzA/1000 row ft.	IF	0.10 ± 0.04f
9	A14776 @ 0.15 OzA/1000 row ft.	IF	0.12 ± 0.05ef
10	Capture 2EC @ 0.33 OzA/1000 row ft.	Band	0.07 ± 0.02f
11	Capture 2EC @ 0.33 OzA/1000 row ft.	IF	0.20 ± 0.08def
12	Aztec 2.1G @ 6.7 OzPr/1000 row ft.	Band	0.01 ± 0.01f
13	Aztec 2.1G @ 6.7 OzPr/1000 row ft.	IF	0.03 ± 0.01f
14	Lorsban 15G @ 8 OzPr/1000 row ft.	Band	0.05 ± 0.02f
15	Lorsban 15G @ 8 OzPr/1000 row ft.	IF	0.06 ± 0.03f
16	Poncho 1250 ST @ 1.25 mg/kernal	ST	0.04 ± 0.02f
17	Cruiser 5FS @ 1.125 gm/100 kg seed	ST	1.05 ± 0.22a
18	YieldGard + Poncho 250 @ 0.25 MgA/seed		0.01 ± 0.01f
19	A14115 FS @ 0.138 GA/100 kg seed Cruiser 5FS @ 1.125 GA/100 kg seed	ST	0.08 ± 0.03f
20	STP15199 @ 5 MgA/seed + STP15201 @ 1.25 MgA/seed	ST	0.02 ± 0.01f
21	Force ST @ 1 MgA/seed	ST	0.26 ± 0.13def
22	A13219 ST @ 1 MgA/seed	ST	0.24 ± 0.12def

23	Cruiser 5FS @1 MgA/seed Force ST @ 1 MgA/seed	ST	0.21 ± 0.05def
24	Cruiser 5FS @1 MgA/seed A13219 ST @ 1 MgA/seed	ST	0.09 ± 0.03f
25	Regent 4 SC @ .24 fl. oz./1000 ft	IF	0.02 ± 0.01f
26	Regent 4SC @ .24 fl. oz./1000 ft + liquid fertilizer	IF	0.03 ± 0.03f

Means within a column followed by the same letter are not significantly different ($P > 0.05$; PROC GLM; Mean comparison by LSD [SAS Institute 2003]).

Soil Insecticide – Test 3 (Yield Data)
 Corn Rootworm – Corn (2005)
 Planting date: April 29, 2005
 Harvest date: September 13, 2005

Gerald Wilde, Department of Entomology, Kansas State University

Trt. No.	Treatment/ Product Name	Application	Yield (bu/acre) Mean ± SE
1	Untreated Check	----	98.44 ± 3.93abcd
2	Force 3G @ 4 OzPr/1000 row ft.	Band	108.75 ± 12.55abc
3	Force 3G @ 4 OzPr/1000 row ft.	IF	107.95 ± 8.75abcd
4	A14776 @ 0.09 OzA/1000 row ft.	Band	83.85 ± 1.31cde
5	A14776 @ 0.12 OzA/1000 row ft.	Band	95.83 ± 7.22abcde
6	A14776 @ 0.15 OzA/1000 row ft.	Band	92.50 ± 5.38abcde
7	A14776 @ 0.09 OzA/1000 row ft.	IF	96.99 ± 7.58abcde
8	A14776 @ 0.12 OzA/1000 row ft.	IF	91.67 ± 13.09abcde
9	A14776 @ 0.15 OzA/1000 row ft.	IF	114.36 ± 12.36a
10	Capture 2EC @ 0.33 OzA/1000 row ft.	Band	90.54 ± 4.70abcde
11	Capture 2EC @ 0.33 OzA/1000 row ft.	IF	87.81 ± 4.63bcde
12	Aztec 2.1G @ 6.7 OzPr/1000 row ft.	Band	95.64 ± 7.58abcde
13	Aztec 2.1G @ 6.7 OzPr/1000 row ft.	IF	105.68 ± 7.55abcd
14	Lorsban 15G @ 8 OzPr/1000 row ft.	Band	78.13 ± 10.67de
15	Lorsban 15G @ 8 OzPr/1000 row ft.	IF	109.69 ± 8.11ab
16	Poncho 1250 ST @ 1.25 mg/kernal	ST	83.04 ± 8.43e
17	Cruiser 5FS @ 1.125 gm/100 kg seed	ST	95.83 ± 18.16abcde
18	YieldGard + Poncho 250 @ 0.25 MgA/seed		92.95 ± 8.22abcde
19	A14115 FS @ 0.138 GA/100 kg seed Cruiser 5FS @ 1.125 GA/100 kg seed	ST	94.69 ± 16.44abcde
20	STP15199 @ 5 MgA/seed + STP15201 @ 1.25 MgA/seed	ST	85.31 ± 5.46bcde
21	Force ST @ 1 MgA/seed	ST	98.96 ± 9.68abcde
22	A13219 ST @ 1 MgA/seed	ST	95.73 ± 9.95abcde

23	Cruiser 5FS @1 MgA/seed Force ST @ 1 MgA/seed	ST	96.25 ± 2.39abcde
24	Cruiser 5FS @1 MgA/seed A13219 ST @ 1 MgA/seed	ST	86.25 ± 1.85bcde
25	Regent 4 SC @ .24 fl. oz./1000 ft	IF	103.13 ± 1.99abcde
26	Regent 4SC @ .24 fl. oz./1000 ft + liquid fertilizer	IF	100.21 ± 11.46abcde

Means within a column followed by the same letter are not significantly different ($P > 0.05$; PROC GLM; Mean comparison by LSD [SAS Institute 2003]).

Reference to specific products is provided solely for informational purposes. Experiments with pesticides on non-labeled crops or pests is part of the insecticide registration process, it does not imply endorsement or recommendation of non-labeled uses of pesticides by Kansas State University. All pesticide use must be consistent with current labels.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating. Fred A. Cholick, Director.