Sunflower Head Moth Control on Sunflower with Foliar Insecticides - 2000.
G. Wilde, Kansas State University.

Pest: Sunflower Moth, *Homeosoma ellectellum*

Crop: Sunflower, DK 3900

Planting Date: April 14

Location: Manhattan, KS

Application: July 5

Treatment: Applied with a single nozzle sprayer at 20 gal/acre at 30 psi. 50% bloom

Evaluated: July 24, by examining 2 head/rep and counting worm/head.

Experimental Design – Randomized Complete Block: 1 row 30 ft. in length, 3 replications.

Sunflower Head Moth Control on Sunflower with Foliar Insecticides - 2000.

Treatment	Form	Al/Acre	Worms/Head
Fury	1.5	0.05	3.2 b
Fury	1.5	0.037	3.3 b
Fury	1.5	0.029	4.2 b
Baythroid	2E	0.043	6.3 b
Baythroid	2E	0.037	9.5 b
Ethyl Parathion	8E	1.0	11.2 b
Asana	0.66	0.03	11.8 b
Warrior	1E	0.03	13.3 b
Lorsban	4E	0.75	14.0 b
Baythroid	2E	0.031	14.2 b
Warrior	1E	0.02	17.5 b
Untreated	n/a	n/a	38.8 a
Thuricide	0.80%	8.6 fl oz	49.3 a

LSD – 15.425

Click here for a PDF file of this Document.

"Knowledge for Life"

Efficacy Trials
Department of Entomology Home Page | Entomology Extension Home page |
K-State Research and Extension Home Page | K-State Home Page |
Entomology Site Map
This page was last updated on: 08/11/04 We hope you enjoy visiting our WWW site.
Please mail any questions, suggestions, or comments to: Psloderb at OzNet.KSU.EDU
