

2018 Alfalfa Insecticide Efficacy Trial – Dickinson Co., KS.

Jeff Whitworth, Holly Davis, JR Ewing - Department of Entomology, Kansas State University

Pest: Alfalfa weevil, Hypera postica, Pea aphid, Acyrthosiphon pisum

Crop: Alfalfa; Established Fall 2017

Location: Dickinson Co., KS

Plot Size: 10 ft. x 30 ft.

Experimental Design: Randomized Complete Block; 4 Replications

Information: Sprayed with hand sprayer delivering 15 gal/acre at ca.30 psi on

27 April, 2018. Plants 2" – 11" tall, temperature 78°F, wind WSW 3-10 mph. Treatments 7 and 9 sprayed again on 4 May 2018.

Temperature 71°F, wind NW 6 mph.

Special Notes: Lowest infestation levels that we've seen in 20+ years due to sub-

freezing temperatures which occurred several times during April. Each sampling period we detected large numbers of very small

dead larvae

Phytotoxicity: None noted

Evaluation: Pre-treatment counts conducted on 28 April. Average of 3.8 larvae/ 10

stems. 10 stems randomly selected in each plot and shaken into 1 gal. bucket and counted 30 April (3 DAT), 4 May (7 DAT), 11 May (14 DAT). At this time alfalfa began flowering and efficacy trial was terminated so that

field could be swathed.

DAT = Days After Treatment

2018 Alfalfa Insecticide Efficacy Trial – Dickinson Co., KS.

Jeff Whitworth, Holly Davis, JR Ewing - Department of Entomology
Kansas State University

Treatment Date: 27 April, 2018

		Mean No. AW Larvae/10 stems			Mean No. Pea	
					Aphids/10 stems	
No.	Treatment	30 April	4 May	11 May	30 April	11 May
		(3 DAT)	(7 DAT)	(14 DAT)	(3 DAT)	(14 DAT)
1	Untreated	4.3a	2.3a	1.3a	2.8a	25.5bcd
2	Endigo ZC 4.5 oz/a	0.0b	0.0c	0.3b	0.0b	18.8cd
3	Warrior II 1.92 oz/a + Lorsban 1 pt/a	0.0b	0.0c	0.0b	0.5ab	14.8cd
4	Warrior II 1.92 oz/a	0.3b	0.0c	0.0b	0.5ab	11.5d
5	Steward EC + silicone surf. ½% 6 oz/a	0.0b	0.0c	0.0b	2.5ab	56.8a
6	Steward EC + silicone surf. ½% 11.3 oz/a	0.0b	0.0c	0.0b	2.0ab	41.0abc
7	Mustang Maxx 4 oz/a followed by Steward EC 8 oz/a -					
	(4 May)	0.5b	0.5b	0.0b	0.8ab	33.8abcd
8	Mustang Maxx 4 oz/a + Dimethoate 4E 16 oz/a	0.0b	0.0c	0.0b	0.0b	18.0cd
9	Steward EC 5.3 oz/a followed by Steward EC 6.0 oz/a			_		_
	(silicone surf. ½%) – (4 May)	0.0b	0.0c	0.0b	2.3ab	39.8abc
10	Steward EC 6.0 oz/a + Mustang Maxx 4 oz/a	0.0b	0.0c	0.0b	0.3ab	47.5ab
11	Besiege @ 8oz/a	0.0b	0.0c	0.0b	0.5ab	25.3bcd

Means within a column followed by the same letter are not significantly different (*P*>0.05; PROC ANOVA; Mean comparison by LSD [SAS Institute 2003]). Reference to specific products is provided solely for informational purposes. Experiments with pesticides on non-labeled crops or pests is part of the insecticide registration process, it does not imply endorsement or recommendation of non-labeled uses of pesticides by Kansas State University. All pesticide use must be consistent with current labels.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, Extension Districts, and United States Department of Agriculture Cooperating, John Floros, Director.